

## EVALUATION OF FAMILY-MANAGED SMALL FARMS USING EMERGY METHODOLOGY

*Agostinho<sup>1</sup>, F.D.R.; Ortega<sup>1</sup>, E.; Diniz<sup>2</sup>, G*

### ABSTRACT

Two small farms, Duas Cachoeiras and Santa Helena, located in the southeast of Brazil (Amparo County, State of São Paulo), were compared using emergy methodology. The first one has been managed using agroecological concepts since 1985, while the second works according with conventional chemical technology. Both are family managed and were formed by the dismemberment of a coffee plantation 30 years ago. The results of this research show that: (a) it is economically feasible to change from a conventional agriculture to agroecological production; (b) the emergy indices show a better performance of the ecological farm; (c) accounting for the partial renewabilities of each incoming flow in the emergy evaluation allowed a more accurate calculation of sustainability; (d) the use of emergy methodology could improve the administration strategies of the farms.

Keywords: Agroecology, Sustainable Development, Emergy, Environment, Ecological Economy.

### 1. INTRODUCTION

The increase of farm productivity in terms of single product yields, due to the Second Agricultural Revolution which took place in the beginning of XX century, led to the replacement of farming systems based on rotation of crops and integration with animal production, by highly specialized monoculture farming systems. This new kind of farms is characterized by massive use of industrial products and fossil energy, as oil, agrochemicals, machinery and vegetal varieties highly responsive to chemical fertilizers. This chemical farming model was reinforced in the 70's by the so called Green Revolution resulting in considerable increase of crop volume, but at the same time, outcoming in environmental and social problems, as forests destruction, biodiversity loss, reduction of water infiltration to aquifers, pollution of water resources, soil erosion, contamination of foods.

In Brazil, besides these problems, it was produced an unfair concentration of lands ownership and wealth distribution, resulting in an intense exodus from countryside to industrialized urban areas [1]. The soil fertility has been diminishing because the adopted techniques are detrimentally affecting soil physical, chemical and biological parameters. Erosion and compacting of soil affect negatively the water supply, both for agricultural and urban use. According to the Food and Agriculture Organization (FAO), the chemical agriculture produces not only food and fiber, but great damages on ecosystems and on local, national and world economies. These impacts are called "negative externalities" and are defined as the costs of the use of the environment for the production of specific product, that currently are not accounted for in the final price of the product itself, encouraging that way conventional models of production [3, 4]. Oil became the basis of industrial agriculture; approximately, 85% of the energy incorporated in foods through farming comes from it [2]. Farming moved to a pattern that can be only sustained as long as fossil fuels are available.

Emergy methodology has being used, since the 90's, as an important tool in studying agricultural systems, in order to suggest a rational exploitation of natural resources [5].

---

<sup>1</sup> Laboratory of Ecological Engineering, Food Engineering School, Campinas State University, Brazil. Correspondence to: Enrique Ortega, Unicamp CP 6121, Campinas, São Paulo, Brasil. 13083-862. Telephone: (19) 37884035; Fax: (19)37884027; E-mail: ortega@fea.unicamp.br

<sup>2</sup> Duas Cachoeiras Farm – [www.sitioduascachoeiras.com.br](http://www.sitioduascachoeiras.com.br)

This research studies, through energy and economic indices, two rural unit using different models of production in order to explain its performance.

## 2. MATERIAL AND METHODS

Two family managed farms were compared using energy indices: (a) Duas Cachoeiras (DC) and (b) Santa Helena (SH). The first one by means of agroecology since 1985, while the second works conventionally. They are located in the in São Paulo, Brazil (Figure 1). The first step in performing the energy evaluation is the identification of the boundaries of the study area. After that, the main components of the system were identified as well as its exchanges with the external environment. A generic systems diagram was drawn using symbolic energy language (Figure 2) and energy indices (Table 1) were then calculated.




Figure 1. (a) Localization of Amparo county in São Paulo State, Brazil; (b) Localization of Duas Cachoeiras farm (DC) and Santa Helena farm (SH) within Amparo county




Figure 2. Energy systems diagram of a generic production system

Table 1. Emery indices used in environmental accounting

<b>Indices</b>	<b>Expression</b>	<b>Signification</b>
Solar transformity (Tr)	Y/Ep	The ratio of the emery of the output divided by the energy of the products
Emery yield ratio (EYR)	Y/F	The ratio of the emery of the output divided by the emery of those inputs that are fed back from outside the system
Emery investment ratio (EIR)	F/I	The ratio of emery of purchased inputs divided by the emery of free inputs
Renewability (%R)	100(R/Y)	The ratio of the renewable inputs divided by the total emery of system
Emery exchange ratio (EER)	Y/[( $\$$ )x(emdollar)]	The ratio of the delivered emery by the system to economy divided by the emery received by the sells of products

R=renewable resources; N=non-renewable resources; I=R+N= inputs from environment; M= materials; S=services; F=M+S= feedback from economic system; Y=F+I=total emery; Ep= energy of products; \$=money input; emdollar in Brazil.

Table 2. Classification of Emery flows considering the partial renewability

<b>Inputs and services</b>	<b>Description</b>
<b>I: Nature contribution</b>	<b>R + N</b>
R: Renewable resources from nature	Rain, materials and services from preserved areas, inputs from air and soil.
N: Nonrenewable resources from nature	Soil, biodiversity, people exclusion.
<b>F: Feedback from economy</b>	<b>F = M + S</b>
<b>M: Materials</b>	<b>M = M<sub>R</sub> + M<sub>N</sub></b>
M <sub>R</sub> : Renewable materials and energy	Renewable materials from natural origin.
M <sub>N</sub> : Nonrenewable materials and energy	Minerals, chemicals, steel, fuel, etc.
<b>S: Services</b>	<b>S = S<sub>R</sub> + S<sub>N</sub></b>
S <sub>R</sub> : Renewable services	Manpower of renewable sources.
S <sub>N</sub> : Nonrenewable services	Other services, taxes, insurance, etc.
<b>Y: Total emery</b>	<b>Y = I + F</b>

The following equations were used to calculate the economic indices:

$$yield = \frac{(input\ money) - (output\ money)}{(output\ money)}$$

$$real\ yield = \frac{(input\ money) - (output\ money + negatives\ externalities)}{(output\ money + negatives\ externalities)}$$

Where:

Input money = money received for the products sales;

Output money = money expenses with purchasing of materials and payment of services;

Negatives externalities = money that would have to be paid for the environmental and social damages (in this paper, the negative externalities were considered 360 US\$/ha/year - extracted by [3, 4] - for the farm that uses a conventional model of agricultural production).

### 3. RESULTS

After visiting the farms and interviewing the owners, it was possible to be acquainted with the managing and energy systems diagram of each farm (Figures 3 and 4) were drawn. These diagrams show the matter and energy flows and point out which are the flows that need to be considered in the energy evaluation. Evaluation Tables were elaborated (Tables 3 and 5) as well as Aggregated Energy Diagrams (Figures 5 and 6).




Figure 3. Energy systems diagram of Duas Cachoeiras farm, year 2003

Soil nitrogen, phosphorous, potassium and calcium were considered as renewable natural resources, since in the DC farm the agroecological management does not require external chemical inputs.

The green manure and the organic matter amendment supply all the minerals required by crops. The plant uptake of soil minerals does not exceed the replenishment of them by biological processes, therefore they were considered renewable in the emergy evaluation Table (Table 3).




Figure 4. Emergy systems diagram of Santa Helena farm, year 2003

On the contrary, Santa Helena farm is managed by a conventional agricultural practice, characterized by a high input of purchased chemicals (Table 5).

Water springs and a river assures the required water supply for the two farms, and, then, it has been considered as renewable resources.

The soil loss was calculated using the revised universal soil loss equation (RUSLE) in a geographical information systems (GIS). This flow was considered as non-renewable natural resource for the two farms, since it exceeded the soil production by the pedogenetic process.

Both, besides vegetable and animal products, groundwater and forest biomass of the two farms have to be considered in the calculation of the energy of the product (Tables 4 and 6), even though they do not have any market price.

Table 3. Emergy flows supporting 1 hectare of land at Duas Cachoeiras farm in 2003

Note	Item	Amount (unit/yr)	Unit	Emergy intensity (seJ/unit)	Reference for seJ/unit	Solar emergy (seJ/yr) $\times 10^{13}$
<b>Renewable Natural Resources (R):</b>						
1	Sun	1.52E+11	J	1	definition	0.0152
2	Rain	6.25E+10	J	3.10E+04	[7]	194
3	Wind	1.51E+10	J	2.45E+03	[7]	3.69
4	Water spring	2.28E+09	J	8.15E+04	[8]	18.6
5	River water	1.09E+08	J	4.28E+05	[8]	4.67
6	Nitrogen	4.89E+02	Kg	4.05E+13	[9]	1980
7	Phosphor	7.69E+01	Kg	2.20E+13	[9]	169
8	Potassium	2.33E+02	Kg	2.92E+12	[9]	67.9
9	Calcium	2.29E+01	Kg	1.00E+12	[9]	2.29
<b>Non-renewable Natural Resources (N):</b>						
10	Soil loss	2.98E+10	J	1.24E+05	[9]	370
<b>Contribution of the Economy (M):</b>						
11	Depreciation	1.23E+02	US\$	3.30E+12	[10]	40.6
12	Fuel	5.29E+07	J	1.11E+05	[9]	0.587
13	Electricity	3.88E+08	J	2.69E+05	[9]	10.4
14	Maintenance	1.29E+01	US\$	3.30E+12	[10]	4.26
<b>Contribution of the Economy (S):</b>						
15	Simple labor	7.00E+01	US\$	3.30E+12	[10]	23.1
16	Family labor	7.00E+01	US\$	3.30E+12	[10]	23.1
17	Maintenance	1.01E+01	US\$	3.30E+12	[10]	3.33
18	Tax	2.24E+00	US\$	3.30E+12	[10]	0.741
19	Private services	1.35E+00	US\$	3.30E+12	[10]	0.444
20	Phone	1.62E+01	US\$	3.30E+12	[10]	5.33
<b>Total Emergy (Y):</b>						<b>2925</b>

Table 4. Total energy produced by the Duas Cachoeiras farm in the year 2003

Product	Production <sup>A</sup> [kg/year]	Caloric value of the product <sup>B</sup> [kcal/kg]	Energy of the product <sup>D</sup> [J/ha.year]
Corn	2400	3500	1.18E+09
Sunflower	710	4750	4.75E+08
Sunflower oil	213	9000	2.70E+08
Sunflower pie	497	2692	1.89E+08
Beans	387	2620	1.43E+08
Pumpkin	280	150	5.92E+06
Cassava	2000	1330	3.75E+08
Potato Candy	500	1140	8.03E+07
Rice	400	3620	2.04E+08
Soy	192	3630	9.82E+07
Vegetables	3900	80	4.40E+07
Fruits	5000	500	3.52E+08
Wool	36	4500	2.28E+07
Wool carpets	24	4500	1.52E+07
Wool clothes	60	4500	3.81E+07
Honey	600	3125	2.64E+08
Propolis	10	5000	7.05E+06
Wax	20	8000	2.26E+07
Water	-	-	1.25E+10 <sup>E</sup>
Biomass	-	-	1.79E+10 <sup>F</sup>

Product	Production <sup>A</sup> [hour/year]	Caloric value of the product <sup>C</sup> [kcal/hour]	Energy of the product [J/ha.year]
Environmental education	4800	146	9.88E+07
Search	600	146	1.23E+07
<b>Total energy produced:</b>			<b>3.43E+10</b>

<sup>A</sup> Data supplied by the proprietor.

<sup>B</sup> The caloric value of the products was extracted by Brazilian Table Food Composition. <http://www.fcf.usp.br/tabela>. Access 20/05/2004.

<sup>C</sup> 3,500kcal/day.person.

<sup>D</sup> Energy [J/ha/year]=production [kg/year]\*caloric value [kcal/kg]\*4,186 [J/kcal]/farm area[29.7ha]

<sup>E</sup> Water infiltration in the soil = 20% of all annual rain ([11], [12], [13], [14] e [15]). Rain in the region is 1,250 mm/year (1.95E+08 liter/year in 15.6 ha). Water infiltrated is 7.43E+07 liter/year.

Water = 7.43E+04 m<sup>3</sup>/year\* (1/29.7 ha)\*(1,000kg/m<sup>3</sup>)\* (5,000J/kg)= 1.25E+10 J/ha/year.

<sup>F</sup> Biomass = 6,870 kJ/m<sup>2</sup>.year\* (10<sup>4</sup>m<sup>2</sup>/ha)\*(7.75ha/29.7 ha)\*(1,000J/1kJ)= 1.79E+10 J/ha/year.

Annual average of liquid productivity of a tropical forest (100 year) = 36,160 [kJ/m<sup>2</sup>.year]. Source: Ecologia: Aventura na Ciência. Editora Globo, pág.8. 1994. The area of recovery of the farm measure 7.75 ha and is 19 years old.




Figure 5. Duas Cachoeiras farm in the year 2003: (a) Energy systems diagram; (b) Percentage of each one of incoming energy flows in relation to the total energy

Table 5. Energy flows supporting 1 ha of land in Santa Helena farm (SH) in the year 2003

N.	Item	Amount (unit/yr)	Unit	Energy intensity (seJ/unit)	Reference for seJ/unit	Solar energy (seJ/yr) $\times 10^{13}$
<b>Renewable Natural Resources (R):</b>						
1	Sun	1.52E+11	J	1	definition	0.0152
2	Rain	6.25E+10	J	3.10E+04	[7]	194
3	Wind	1.51E+10	J	2.45E+03	[7]	3.69
4	River water	1.25E+09	J	4,28E+05	[8]	53.4
<b>Non-renewable Natural Resources (N):</b>						
5	Soil loss	5.33E+10	J	1.24E+05	[9]	661
<b>Contribution of the Economy (M):</b>						
6	Depreciation	4.77E+02	US\$	3.30E+12	[10]	157
7	Fuel	2.29E+08	J	1.11E+05	[9]	2.55
8	Electricity	1.73E+09	J	2.69E+05	[9]	46.6
9	Materials of maintains	4.27E+01	US\$	3.30E+12	[10]	14.1
10	Fungicide	3.33E+01	Kg	2.49E+13	[9]	82.8
11	Herbicide	4.98E-01	Kg	2.49E+13	[9]	1.24
12	Calcium	2.40E-01	Kg	1.00E+12	[9]	0.0240
13	Calcium nitrate	1.58E+00	US\$	3.30E+12	[10]	0.522
14	Potassium nitrate	2.14E+00	US\$	3.30E+12	[10]	0.705
<b>Contribution of the Economy (S):</b>						
15	Family labor	2.67E+02	US\$	3.30E+12	[10]	88.0
16	Tax	2.46E+00	US\$	3.30E+12	[10]	0.811
17	Private service	8.33E+01	US\$	3.30E+12	[10]	27.5
18	Phone	1.54E+01	US\$	3.30E+12	[10]	5.08
<b>Total Energy (Y):</b>						<b>1340</b>

Table 6. Total energy produced by the Santa Helena farm in the year 2003

Product	Production <sup>A</sup> [kg/year]	Caloric value of the product <sup>B</sup> [kcal/kg]	Energy of the product <sup>C</sup> [J/ha.year]
Coffee	8000	1660	3.56E+09
Chayote	66000	160	2.83E+09
Tomato	9000	150	3.62E+08
Peach	350	410	3.85E+07
Pimienta	2750	80	5.90E+07
Cucumber	11000	60	1.77E+08
Water	-	-	6.25E+09 <sup>D</sup>
Biomass	-	-	5.57E+08 <sup>E</sup>
<b>Total energy produced:</b>			<b>1.38E+10</b>

<sup>A</sup> Data supplied by the proprietor.

<sup>B</sup> The caloric value of the products was extracted by Brazilian Table Food Composition. <http://www.fcf.usp.br/tabela>. Last Access: 20/05/2004.

<sup>C</sup> Energy [J/ha/yr]=production [kg/year]\*caloric value [kcal/kg]\*4186 [J/kcal]/farm area [15.6 ha]

<sup>D</sup> Water infiltration in the soil = 10% of all annual rain in compacted soil ([11], [12], [13], [14] and [15]). Rain in the region is 1250 mm/year (1.95E+08 liter/year in 15.6 ha). The value of water infiltration is 1.95+E07 liter/year (1.95E+04 m<sup>3</sup>/year).

Water = 1.95E+04 m<sup>3</sup>/year \* (1/15.6 ha)\*(1000 kg/m<sup>3</sup>)\* (5000 J/kg)= 6.25E+09 J/ha/year.

<sup>E</sup> Biomass = 1447 kJ/m<sup>2</sup>.year \* (10<sup>4</sup>m<sup>2</sup>/ha)\*(0.6 ha/15.6 ha)\*(1000J/1kJ)= 5.57E+08 J/ha/year.

Annual average of liquid productivity of a tropical forest (100 year) = 36160 [kJ/m<sup>2</sup>.year]. Source: Ecologia: Aventura na Ciência. Editora Globo. Page 8. 1994.

The forest recovery area of this farm measures 0.6 ha and is 4 years old.




Figure 6. Santa Helena farm in the year 2003: (a) Aggregated energy systems diagram; (b) Percentage of the incoming energy flows in relation to the total energy

In Tables 7 and 9 the farm energy evaluations has been also carried out considering the partial renewabilities of each incoming flow in the system.

Table 7. Energy flows supporting 1 hectare of land in Duas Cachoeiras farm (DC) in the year 2003, accounting for the partial renewabilities of each component

N.	Item	Ren. n. <sup>A</sup>	Numerical Value <sup>B</sup>	Unit	Energy intensity <sup>B</sup> [seJ/Unit.]	Renew. Energy <sup>C</sup>	Non- renew. Energy <sup>C</sup>	Y <sup>C</sup>
<b>Renewable Natural Resources (R):</b>								
1	Sun	1	1.52E+11	J	1	0.0152	0.0	0.015
2	Rain	1	6.25E+10	J	3.10E+04	194	0.0	2
3	Wind	1	1.51E+10	J	2.45E+03	3.69	0.0	194
4	Water spring	1	2.28E+09	J	8.15E+04	18.6	0.0	3.69
5	River water	1	1.09E+08	J	4.28E+05	4.67	0.0	18.6
6	nitrogen	1	4.89E+02	kg	4.05E+13	1980	0.0	4.67
7	Phosphorous	1	7.69E+01	kg	2.20E+13	169	0.0	1980
8	Potassium	1	2.33E+02	kg	2.92E+12	67.9	0.0	169
9	Calcium	1	2.29E+01	kg	1.00E+12	2.29	0.0	67.9
<b>Non-renewable Natural Resources (N):</b>								
10	Soil loss	0	2.98E+10	J	1.24E+05	0.0	370	370
<b>Contribution of the Economy (M):</b>								
11	Depreciation	0.0 5	1.23E+02	US\$	3.30E+12	2.03	38.6	40.6
12	Fuel	0	5.29E+07	J	1.11E+05	0.0	0.587	0.587
13	Electricity	0.7	3.88E+08	J	2.69E+05	7.30	3.13	10.4
14	Materials	0.1	1.29E+01	US\$	3.30E+12	0.426	3.83	4.26
<b>Contribution of the Economy (S):</b>								
15	Simple labor	0.6	7.00E+01	US\$	3.30E+12	13.9	9.24	23.1
16	Family labor	0.9	7.00E+01	US\$	3.30E+12	20.8	2.31	23.1
17	Maintenance	0.1	1.01E+01	US\$	3.30E+12	0.333	3.00	3.33
18	Tax	0.0 5	2.24E+00	US\$	3.30E+12	0.0370	0.704	0.741
19	Service	0.0 5	1.35E+00	US\$	3.30E+12	0.0222	0.422	0.444
20	Phone	0.0 5	1.62E+01	US\$	3.30E+12	0.267	5.07	5.33
<b>Total Energy (Y):</b>								<b>2925</b>

<sup>A</sup> Extract from [6] and common sense.

<sup>B</sup> The numeric value and energy intensity are the same as in Table 3.

<sup>C</sup> These flows are multiplied by  $10^{13}$  [seJ/ha/year].

Table 8. Aggregate flows of Duas Cachoeiras farm (DC) considering partial renewabilities in the year 2003

Renewable <sup>A</sup>			Non-Renewable <sup>A</sup>		
Rr =	2.443E+16	seJ/ha/year	Rn =	0.0	seJ/ha/year
Nr =	0.0	seJ/ha/year	Nn =	3.70E+15	seJ/ha/year
Mr =	9.760E+13	seJ/ha/year	Mn =	4.61E+14	seJ/ha/year
Sr =	3.533E+14	seJ/ha/year	Sn =	2.07E+14	seJ/ha/year

<sup>A</sup> Data come from of Table 7.

The next step was the calculation of energy and economic indices (Tables 11 and 12).

Table 9. Emergy flows supporting 1 hectare of land in Santa Helena farm in the year 2003, accounting for the partial renewabilities of each component

N.	Item	R <sup>A</sup>	Numerical Value <sup>B</sup>	Unit	Emergy intensity <sup>B</sup> [seJ/Unit.]	Ren. Em. <sup>C</sup>	Non-ren. Em. <sup>C</sup>	Y <sup>C</sup>
<b>Renewable Natural Resources (R):</b>								
1	Sun	1	1.52E+11	J	1	0.0152	0.0	0.0152
2	Rain	1	6.25E+10	J	3.10E+04	194	0.0	194
3	Wind	1	1.51E+10	J	2.45E+03	3.69	0.0	3.69
4	River water	1	1.25E+09	J	4.28E+05	53.4	0.0	53.4
<b>Non-Renewable Natural Resources (N):</b>								
5	Soil loss	0	5.33E+10	J	1.24E+05	0.0	661	661
<b>Contribution of the Economy (M):</b>								
6	Depreciation	0.05	4.77E+02	US\$	3.30E+12	7.87	150	157
7	Fuel	0	2.29E+08	J	1.11E+05	0.0	2.55	2.55
8	Electricity	0.7	1.73E+09	J	2.69E+05	32.6	14.0	46.6
9	Materials	0.1	4.27E+01	US\$	3.30E+12	1.41	12.7	14.1
10	Fungicide	0.05	3.33E+01	kg	2.49E+13	4.14	78.7	82.8
11	Herbicide	0.05	4.98E-01	kg	2.49E+13	0.0620	1.18	1.24
12	Calcium	0.05	2.40E-01	kg	1.00E+12	0.0012	0.0228	0.0240
13	Nitr. Calcium	0.05	1.58E+00	US\$	3.30E+12	0.0261	0.496	0.522
14	Nitr. Potass.	0.05	2.14E+00	US\$	3.30E+12	0.0353	0.670	0.705
<b>Contribution of the Economy (S):</b>								
15	Family labor	0.9	2.67E+02	US\$	3.30E+12	79.2	8.80	88.0
16	Tax	0.05	2.46E+00	US\$	3.30E+12	0.0405	0.770	0.811
17	Service	0.05	8.33E+01	US\$	3.30E+12	1.38	26.1	27.5
18	Phone	0.05	1.54E+01	US\$	3.30E+12	0.254	4.82	5.08
<b>Total Emergy (Y):</b>								<b>1340</b>

<sup>A</sup> Extract from [6] and common sense.

<sup>B</sup> The numeric value and emergy intensity are the same as in Table 5.

<sup>C</sup> These flows are multiplied by 10<sup>13</sup> [seJ/ha/year].

Table 10. Aggregate flows of the Santa Helena farm (SH) considering partial renewabilities in the year 2003.

Renewable <sup>A</sup>			Non-Renewable <sup>A</sup>		
Rr =	2.508E+15	seJ/ha/year	Rn =	0.0	seJ/ha/year
Nr =	0.0	seJ/ha/year	Nn =	6.61E+15	seJ/ha/year
Mr =	4.614E+14	seJ/ha/year	Mn =	2.60E+15	seJ/ha/year
Sr =	8.087E+14	seJ/ha/year	Sn =	4.05E+14	seJ/ha/year

<sup>A</sup> Data come from Table 9.

Table 11. Emergy indices without considering and considering the partial renewabilities.

Indices (traditional)	Indices (+renewabilities)	traditional		+renewabilities	
		DC <sup>A</sup>	SH <sup>A</sup>	DC	SH
Tr = Y/Ep [seJ/J]	Tr=Y/Ep [seJ/J]	851,7	968,0	851,7	968,0
EYR=Y/F	EYR=Y/(Mn+Sn)	26.12	3.13	43.73	4.46
EIR=F/I	EIR=(Mn+Sn)/(R+Mr+Sr+N)	0.04	0.47	0.02	0.29
%R=100*(R/Y)	%R=100*(R+Mr+Sr)/Y	83.52	18.72	85.06	28.20
EER=Y/input.emdollar <sup>B</sup>	EER=Y/input.emdollar <sup>B</sup>	11.79	2.64	11.79	2.64

<sup>A</sup> DC = Duas Cachoeiras farm in the year 2003; SH = Santa Helena farm in the year 2003

<sup>B</sup> emdollar in Brazil (2003) = 3.3 10<sup>12</sup> seJ/US\$. Reference: [10]  
input (money): DC = 67 000 R\$/year; SH = 71 890 R\$/year

Table 12. Economic Indices

Indices	Equation	DC <sup>A</sup>	SH <sup>A</sup>
Yield <sup>B</sup>	$\frac{(\text{input money}) - (\text{output money})}{(\text{output money})}$	1.93	1.11
Real Yield <sup>C</sup>	$\frac{(\text{input money}) - (\text{output money} + \text{negative externalities})}{(\text{output money} + \text{negative externalities})}$	1.93	0.41

<sup>A</sup> DC = Duas Cachoeiras farm in the year 2003; SH = Santa Helena farm in the year 2003.

<sup>B</sup> output money: DC = 22 845 R\$/year; SH = 34 093 R\$/year.

<sup>C</sup> In this calculation the value of the negative externalities was only considered in conventional production. The considered value was 360 US\$/ha/year [3, 4].

## 4. DISCUSSION

### 4.1. Transformity

The transformity of Duas Cachoeiras farm (851,000 seJ/J) is 12% lower than Santa Helena farm value (967,000 seJ/J), pointing out a best ecosystem efficiency of DC farm (Table 11), due to the diversification and recycling, which are characteristic of ecological farming. The low values of both transformities can be explained by the small dimension of both farms and the intense use of human labor, mostly provided by family members.

### 4.2. Emergy Yield Ratio (EYR)

The EYR is an indicator that measures the ability to exploit local natural energy resources, both renewable and non-renewable. For Duas Cachoeiras farm EYR is 26.1, whereas for Santa Helena farm the value was 3.1 (Table 11). This result points out that the first farm uses much more natural resources, showing less dependence from non-renewable resources. Intensive conventional agricultural systems have EYR values lower of than 1.1 [16].

### 4.3. Emergy Investment Ratio (EIR)

EIR measures the ratio between the investment of society and the contribution of nature to produce a specific good. The EIR index accounted 0.04 for the Duas Cachoeiras farm and 0.47 for the Santa Helena farm. Thus, for the Duas Cachoeiras farm, the environment provides more resources used in the productive process, as compared to the Santa Helena farm. This situation should lower the production costs allowing a better performance on the market. This index accounts, for highly conventional agricultural systems, about 7 [16].

### 4.4. Renewability (%R)

The renewability observed in this study was 83 and 18% for the Duas Cachoeiras farm and the Santa Helena farm, respectively (Table 11). The agroecological management is closer to the self-sustainability, with higher quality of the products, issuing in a higher price on the market, and lower dependence on external resources. The recommendations towards a more sustainable agricultural system, suggested by the Agenda 21, at global and national levels, could lead to adjustments on farms producing conventionally, reducing environmental and social impacts. The use of partial renewability of each input used lead to a better measurement of renewability. Accounting for the renewability of each input can improve emergy evaluations, it is especially for important for semi-renewable inputs (as manure, labor, electricity) purchased at local or regional scales of the economy [6]. The DC index changes from 83% to 85%, while the SH changes from 18% to 28%. There is an important difference in the renewable fraction of family labor

(90% of renewability) and contracted labor (60% of renewability). These farms use family labor or employees living in the same region where the property is located. In conventional systems of production, where all labor force is made by hired workers, these flows are considered not renewable [6].

#### **4.5. Emergy Exchange Ratio (EER)**

This index verifies if the total emergy used in the system was paid (in monetary terms) by selling the products. The lower this value the better, since it is calculated by dividing the emergy used by the emergy received. The EER obtained were 11.79 for the Duas Cachoeiras farm and 2.64 for the Santa Helena farm (Table 11), indicating a better performance of Santa Helena farm.

#### **4.6. Economic yield**

The income calculated in this research was 193% for Duas Cachoeiras farm and 111% Santa Helena farm (Table 12). Thus, for each monetary unit spent in the Duas Cachoeiras farm, 2.9 monetary units are earned by products selling, while for the Santa Helena farm, 2.1 monetary units are earned. This result points out that both properties obtain profit. Duas Cachoeiras farm, however, obtains higher profit, protecting nature in the meanwhile. Considering the costs of the negative externalities in the calculation of the income for conventional system (Table 12), the index decreases of 41% for the Santa Helena farm, while in the Duas Cahoeiras farm, the index does not change, because the farm does not produce any negative externality.

### **5. CONCLUSIONS**

The quality of farm diagnosis obtained by the interpretation of emergy indices confirms that emergy methodology is useful tool in performing environmental accounting of production systems, since it takes into consideration the contribution of the nature beyond production means, labor and services. The application of emergy analysis could improve the development of planning and administration tools needed to achieve a more sustainable development, following Agenda 21 commitments.

This research leads to the following conclusions: (a) emergy and economic indices showed a better performance for the Duas Cachoeiras farm, indicating that the use of agroecology results in a more balanced relationship between production and natural environment; (b) since both farms are located in the same region and have the same physical characteristics (soil, altitude, climate, solar radiation, rain, regional biodiversity, etc.), the differences in emergy and monetary values between them are only due to the agricultural model adopted; (c) accounting for the partial renewability of each incoming flow allowed a more accurate measurement of the sustainability of the farms under study; (d) emergy methodology could allow a better farm management resulting in higher efficiency and an environmentally friendly approach.

### **ACKNOWLEDGEMENTS**

The authors are grateful to Antonio Padilha for his aid in collecting data used in this paper, and to Consuelo Pereira and Otávio Cavallet for their contribution to this research.

## References

- [01] Ehlers, E.; Agricultura sustentável – Origens e Perspectivas de um Novo Paradigma. Ed. Livros da Terra. São Paulo. 1996.
- [02] Ulgiati, S.; Odum, H.T.; Bastianoni, S. Emergy use, environmental loading and sustainability: in emergy analysis of Italy. *Ecological Modelling*. V.73, p.215-268. 1994.
- [03] Pretty, J.N.; Brett, C.; Gee, D.; Hine, R.E.; Mason, C.F.; Morison, J.I.L.; Raven, H.; Rayment, M.D.; van der Bijl, G; An assessment of the total external costs of UK agriculture. *ELSEVIER - Agricultural Systems*, 65, 113-136. 2000.
- [04] Pretty, J.N.; Brett, C.; Gee, D.; Hine, R.E.; Mason, C.F.; Morison, J.I.L.; Rayment, M.D.; van der Bijl, G; Dobbs, T.; POLICY AND PRACTICE: Policy Challenges and Priorities for Internalizing the Externalities of Modern Agriculture. *Journal of Environmental Planning and Management*, 44(2), 263-283. 2001
- [05] Comar, M.V.; Avaliação emergética de projetos agrícolas e agro-industriais no Alto Rio Pardo: a busca do Desenvolvimento Sustentável. Tese de Doutorado. Campinas-UNICAMP. 1998.
- [06] Ortega, E.; Anami, M.; Diniz, G.; Certification of food products using emergy analysis. In: *Proceedings of III International Workshop Advances in Energy Studies: reconsidering the importance of energy*. September, 24-28, Porto Venere, Italy, p. 227-237. 2002.
- [07] Odum, H.T., Brown, M.T. and Brandt-Williams, S. L., 2000. *Handbook of Emergy Evaluation: A Compendium of Data for Emergy Computation Issued in a Series of Folios. Folio No.1 - Introduction and Global Budget*. Center for Environmental Policy, Environmental Engineering Sciences, Univ. of Florida, Gainesville, 17 pp. Available in: <http://www.ees.ufl.edu/cep/> Access in 10 oct. 04.
- [08] Bastianoni, S. and Marchettini, N., 2000. The problem of co-production in environmental accounting by emergy analysis. *Ecol. Model.* 129: 187-193.
- [09] Brandt-Williams, S. L., 2002. *Handbook of Emergy Evaluation: A Compendium of Data for Emergy Computation Issued in a Series of Folios. Folio No.4 - Emergy of Florida Agriculture*. Center for Environmental Policy, Environmental Engineering Sciences, Univ. of Florida, Gainesville, 40 pp. Available in: <http://www.ees.ufl.edu/cep/> Access in 10 oct. 04.
- [10] Coelho, O., Ortega, E. & Comar, V., 2003. Balanço de Emergia do Brasil (Dados de 1996, 1989 e 1981). (Emergy balance of Brazil - Statistics of 1996, 1989 e 1981). In: *Engenharia Ecológica e Agricultura Sustentável (Ecological Engineering and Sustainable Agriculture)*. Organizer: Enrique Ortega. Available in: <http://www.fea.unicamp.br/docentes/ortega/livro/index.htm> Access in: 15/07/2004.
- [11] Cavenage, A.; Moraes, M.L.T.; Alves, M.C.A.; Carvalho, M.A.C.; Freitas, M.L.M.; Buzetti, S.; Alterações nas propriedades físicas de um Latossolo Vermelho escuro, sob diferentes culturas. *Revista Brasileira de Ciência do Solo*. v.23, nº.4, p.997-1003. 1999.
- [12] Centurion, J.F.; Cardoso, J.P.; Natale, W.; Efeito de formas de manejo em algumas propriedades físicas e químicas de um Latossolo Vermelho em diferentes agroecossistemas. *Revista Brasileira de Engenharia Agrícola e Ambiental*. v.5, nº.2, p.254-258. 2001.
- [13] Gicheru, P.; Gachene, C.; Mbuvi, J.; Mare, E. ; Effects of soil management practices and tillage systems on surface soil water conservation and formation on a sandy loam in semi-arid Kenya. *Soil & Tillage Research*. 75: 173-184. 2004.
- [14] Lima, W. de P.; Impacto ambiental do eucalipto. 2ª edição. São Paulo: Editora da Universidade de São Paulo. 1996.
- [15] Souza, Z.M. de; Alvez, M.C.; Movimento de água e resistência à penetração em um Latossolo Vermelho distrófico de cerrado, sob diferentes usos e manejos. *Revista Brasileira de Engenharia Agrícola e Ambiental*. v.7, nº.1, p.18-23. 2003.
- [16] Odum, H.T.; *Environmental Accounting: Emergy and Environmental Decision Making*. John Wiley & Sons, Inc., New York, USA. 1996.