

How Brazil is dealing with COVID-19 pandemic arrival regarding elective gynecological surgeries

Luiz Gustavo Oliveira Brito MD/PhD ,  
Paulo Ayrosa Ribeiro MD/PhD ,  
Agnaldo Lopes Silva-Filho MD/PhD , FEBRASGO Brazilian  
Gynecological Surgery Group for COVID-19

PII: S1553-4650(20)30216-8  
DOI: <https://doi.org/10.1016/j.jmig.2020.04.028>  
Reference: JMIG 4128

To appear in: *The Journal of Minimally Invasive Gynecology*

Received date: 20 April 2020  
Accepted date: 22 April 2020

Please cite this article as: Luiz Gustavo Oliveira Brito MD/PhD , Paulo Ayrosa Ribeiro MD/PhD , Agnaldo Lopes Silva-Filho MD/PhD , FEBRASGO Brazilian Gynecological Surgery Group for COVID-19, How Brazil is dealing with COVID-19 pandemic arrival regarding elective gynecological surgeries, *The Journal of Minimally Invasive Gynecology* (2020), doi: <https://doi.org/10.1016/j.jmig.2020.04.028>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier Inc. on behalf of AAGL.



**How Brazil is dealing with COVID-19 pandemic arrival regarding elective gynecological surgeries**

**Luiz Gustavo Oliveira Brito, MD/PhD<sup>1,4</sup>,**

**Paulo Ayrosa Ribeiro, MD/PhD<sup>2,4</sup>**

**Agnaldo Lopes Silva-Filho, MD/PhD<sup>3,4</sup> and**

**FEBRASGO Brazilian Gynecological Surgery Group for COVID-19<sup>\*4</sup>**

**1** – Department of Obstetrics and Gynecology, University of Campinas, Campinas, Brazil

**2** – Department of Obstetrics and Gynecology, Santa Casa School of Medical Sciences, São Paulo, Brazil

**3** – Department of Gynecology and Obstetrics, Federal University of Minas Gerais, Belo Horizonte, Brazil

**4** – Executive Secretariat, Avenida Brigadeiro Luis Antonio, 3421, sala 903 – São Paulo, Brazil

**Corresponding author:**

Luiz Gustavo Oliveira Brito, MD, MsC, PhD

Rua Alexander Fleming, 101 – Cidade Universitária – Campinas - Brazil

Zipcode 13083-881 – Phone: +55-19-3521-9595 – E-mail: lgo Brito@gmail.com

**Manucript word count:** 487 words

**Funding:** none

**Conflict of interests:** none

**Keywords:** COVID-19; Brazil; gynecologic surgical procedures

Brazil, a continental country with 210 million people, is starting to have an increase of infected cases and mortality due to COVID-19 that has started on February (1). Whereas the pandemic has spread worldwide, and our curve is weeks behind other epicenters such as United States, Italy and Spain, our main ObGyn Society (Brazilian Federation of Associations of Gynecology and Obstetrics - FEBRASGO) and several affiliated societies (eg. UROGINAP, SBE) have been preparing to counsel our members.

Several nonpharmaceutical interventions were oriented nationally (2) and discussions between healthcare providers and decision-makers have helped several hospitals (3) prior to the explosion of cases. Obstetrical and oncology services had a smaller reduction of cases than services who are mainly referred for elective surgeries. We have prepared documents in FAQ-format for counseling gynecological surgeons to reschedule elective surgeries for benign diseases; (4) discussed to implement telemedicine in many outpatient clinics and analyzed to reorganize surgical scenarios for gynecologic oncology patients and/or emergency surgeries. However, we are still facing many regional barriers such as: lack of massive testing for patients, either asymptomatic or symptomatic, disparities between the number of COVID-19 patients interned in public intensive care units (ICUs) versus private ICUs (and also number of available ICUs throughout the country), possible undernotification of a central national data for several reasons and most importantly, not all facilities have sufficient personal protective equipment (PPE).

Our guidelines are aligned with the joint statement previously published in AAGL with several societies (5): elective surgeries should be postponed or rescheduled, and this should be a joint decision between surgeon and patient; risk factors for COVID-19 should be checked with patient, and the absence of COVID-19 symptoms should exhaustively be confirmed in case she needs a surgery to undergo.

Testing should be massively encouraged prior to surgery. Within the operating room, N-95 masks should be guaranteed for the surgical team (as well as the rest of accessories – shoe covers, gown, protective head covering, gloves and eye protection), number of personnel should be restricted inside the operating room (OR), and prior conversation with the anesthesia team should be done prior to each procedure.

For vaginal and abdominal surgeries, use of N-95 masks plus face shields should be present and smoke dispersion should be avoided as much as possible. Regarding laparoscopic procedures, it can be performed after informed consent about the subject, if the entire OR team has access to necessary PPE and extreme care is taken to reduce pneumoperitoneum escape (6). It is important to remember that to this moment, no data are available proving that COVID-19 viral particles were identified in surgical smoke. Finally, most urogynecological disorders can be treated conservatively.

We are still uncertain whether we are flattening the curve, and discussions about reopening elective surgeries are still controversial; however, we believe we are ready to fight this battle and we are extensively advising patients to seek hospitals if their symptoms are increasing during the quarantine.

***\*This group is composed by:*** Mariano Tamura, MD, PhD; Julio Cesar Rosa-e-Silva, MD, PhD; Ricardo Quinteiros, MD; Helizabet Abdalla-Ribeiro MD, PhD; Sergio Podgaec, MD, PhD; Walquiria Primo, MD; Carlos Augusto Lino, MD; Mauricio S. Abrao, MD, PhD; Renato Moretti-Marques, MD, PhD

## References

1 – Rodriguez-Morales AJ, Gallego V, Escalera-Antezana JP et al. COVID-19 in Latin America: the implications of the first confirmed case in Brazil. *Travel Med Infect Dis* 2020;29:101613.

2 – Garcia LP, Duarte E. Nonpharmaceutical interventions for tackling the COVID-19 epidemic in Brazil. *Epidemiol Serv Saude* 2020;29(2):e2020222.

3 – Chen AT, Moniz CM, Ribeiro-Junior U et al. How should health systems prepare for the evolving COVID-19 pandemic? Reflections from the perspective of a tertiary cancer center. *Clinics* 2020;75:e1864.

4 – FEBRASGO. Covid 19. Accessed in Apr 20, 2020. <https://www.febrasgo.org.br/pt/covid19>

5 – AAGL. Joint Statement on Minimally Invasive Gynecologic Surgery. Accessed in : Apr 20, 04, 2020 - <https://www.aagl.org/news/covid-19-joint-statement-on-minimally-invasive-gynecologic-surgery/>

6 – Cohen SL, Liu G, Abrao M, Smart N, Heniford T. Perspectives on surgery in the time of COVID-19: safety first. *J Minim Invasive Gynecol* 2020Apr 3. doi: 10.1016/j.jmig.2020.04.003